


Accepted Manuscript

Title: The Next Generation of Mindfulness-Based Intervention
Research: What have we learned and where are we headed?

Authors: Melissa A Rosenkranz, John D Dunne, Richard J Davidson



PII: S2352-250X(18)30189-1

DOI: <https://doi.org/10.1016/j.copsyc.2018.12.022>

Reference: COPSYC 765

To appear in:

Please cite this article as: Rosenkranz MA, Dunne JD, Davidson RJ, The Next Generation of Mindfulness-Based Intervention Research: What have we learned and where are we headed?, *Current Opinion in Psychology* (2018), <https://doi.org/10.1016/j.copsyc.2018.12.022>

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

The Next Generation of Mindfulness-Based Intervention Research: What have we learned and where are we headed?

Melissa A. Rosenkranz, John D. Dunne, Richard J. Davidson
Center for Healthy Minds, University of Wisconsin-Madison

Abstract

The previous two decades have seen an exponential increase in the number of published scientific investigations on the efficacy of mindfulness-based stress reduction (MBSR) training to improve function in a wide range of physical and psychological processes. The resulting body of work provides strong evidence that MBSR has salubrious effects. Yet, when compared directly to groups with training that matches MBSR in factors common to most legitimate interventions, such as learning new skills, expectation of benefit, social engagement and support, and attention from expert instructors, both groups tend to improve to a similar extent. This raises the question of whether there are benefits that are specific to training in mindfulness and if so, why are we not detecting them? Here, we discuss the factors that contribute to the general lack of differentiation between MBSR and active control groups, including the specificity of outcome measures and experimental design, random assignment, inclusion/exclusion criteria, and the time course and trajectory of change. In addition, we offer recommendations to address these factors in future research.

Since its inception in 1979, Mindfulness-based Stress Reduction (MBSR) training has demonstrated efficacy in promoting health and wellbeing across a broad range of outcome measures^{1,2}. As such, MBSR has become the most widely implemented meditation training program in healthcare settings in the United States, as well as in many other countries around the world. Yet, when directly compared to active control (AC) interventions¹, such as exercise or relaxation, a lack of group differentiation is common^{3–7}. In fact, the frequency of a null result in this regard is likely underestimated, due to a known publication bias in this area⁸. In this article, we discuss a range of possible factors that may contribute to a lack of divergence between MBSR and active control groups. These factors comprise three categories: the specificity of the therapeutic mechanism; experimental design; and the specificity of outcome measures and their analysis. In addition, we offer recommendations to address these factors in future research.

The purpose of an AC is to match the target intervention on common factors (as well as structure, duration, and intensity), as they are important therapeutic components of any bona fide intervention, but are not specific to the content of a particular intervention. Common factors include therapeutic alliance, expectation and rationale for benefit, social support and group cohesion, attention from content experts, and learning new skills and habits⁹. The relationship between the provider and the recipient, or therapeutic alliance, for

¹ It is important to note that the vast majority of studies that examine the efficacy of MBSR employ a wait-list control group, rather than an active control group.

example, alone accounts for more than 7% of the variance in psychotherapy outcome measures¹⁰. In drug trials, this effect is much larger, where expectation of benefit and therapeutic alliance can account for 32% and 22%, respectively, of variance in symptom reduction¹¹. It is important to note that when MBSR is compared to an AC with a null outcome, this result is often due to commensurate post-intervention improvements in both groups, rather than an absence of change. This suggests that improvements shown in both MBSR and ACs may reflect benefits that can be attributed to these common factors. On the contrary, wait-list control conditions lack common factors entirely, and treatment-as-usual (TAU) control groups lack the common factors associated with the bonus intervention (i.e. TAU + MBSR).

A second, closely related factor is that different mechanisms of change can lead to the same outcome. For example, both mindfulness training and aerobic exercise have reliably been shown to improve mood¹²⁻¹⁴ and sleep quality^{15,16}, though the mechanisms through which they do so are likely to be quite different. Among the proposed mechanisms through which mindfulness training improves sleep quality are improved regulation of autonomic arousal, decreased worry and rumination, and increased acceptance^{15,17} via strengthening of meta-awareness and dereification¹⁸, whereas the proposed mechanisms through which aerobic exercise improves sleep quality include effects on thermoregulatory processes, cytokine release, products of metabolism on sleep architecture, and shifts in circadian rhythm^{19,20}. The apparent similarity in outcomes could simply reflect the reality that there are many ways to reach the

same endpoint. Alternatively, it could indicate that overly gross or distal measures are poorly suited to detect fine-grained distinctions in what are actually different endpoints, produced by different mechanisms.

Though some outcome measures, like sleep quality, may not differentiate these interventions regardless of the measurement time frame, for other outcomes, time frame is important. It is not uncommon, for example, for outcomes that *should* be specific to one intervention, such as ability to sustain attention (mindfulness) or lactic acid threshold (aerobic exercise) to show little change or change that is highly variable in the early stages of training^{21,22}. However, after a period of prolonged or intensive training, differentiation in these intervention-specific outcomes becomes apparent^{23,24}. Without the proper study design and clear hypotheses about which changes are likely to occur within a given time frame, one cannot determine whether a null finding resulted from the intervention and AC leading to the same outcome, or whether the measurement interval was insufficient to detect long-term differentiation in outcomes. To address this issue, careful selection of outcome measures is required, with consideration given to the changes that can be realistically expected, in a majority of participants, in response to a particular “dose” of the intervention. In some circumstances, this is an entirely empirical question. In other cases, much insight can be gained through consultation and collaboration with meditation experts. Further, as is common practice in pharmacology, examination of outcomes at different “doses”, i.e. varying the span and intensity of training, is necessary to understand which outcomes *tend* to emerge after a certain amount

of training, and what the appropriate dose may be, given a particular desired outcome².

In addition to the careful matching of outcome measures to the appropriate intervention dose or snapshot in time, the specific relevance of outcome measures to the skills trained in MBSR is critical. For the most part, outcome measures have been drawn from existing areas of inquiry and have not been developed to assess the specific changes we expect to result from this training. There are good reasons behind this approach. First, the ability to translate benefits of the training to more general aspects of wellbeing is highly desirable. Second, choosing measures from another area of inquiry allows one to interpret the outcome in reference to a much larger body of scientific work and to compare the outcomes of mindfulness training to the outcomes from other types of interventions. Finally, development of new tasks is time consuming and costly. Use of existing tasks and surveys is an efficient way to generate data, especially in a nascent field during a period of increasingly tight grant cycle timelines. However, generalizability of skills learned in the training context to other scenarios requires a certain level of mastery, and can vary considerably between individuals. After only 8 weeks of training, this generalization may be minimal, resulting in a situation where MBSR and ACs perform equally. To address this issue, a collaborative effort is needed, where tasks are developed and validated that assess the specific changes expected to follow from a given duration or

² As is discussed in more detail below, change resulting from meditation practice may not be linear and the trajectory of change can vary widely between individuals.

intensity of each type of contemplative practice. For example, in the early stages of training, one might expect to see improvement in skills explicitly trained during the intervention, such as focus on the breath, in a breath-counting task²⁵.

Whereas after prolonged training, generalization of explicitly trained skills to other contexts would be expected, for example in tasks that assess mind-wandering or the ability to sustain attention. This collaborative effort should result in a widely available set of tasks, akin to the NIH Toolbox for assessing cognitive function.

In treatment research, the randomized clinical trial (RCT) is the gold-standard experimental design. This approach was developed primarily in the context of pharmacological intervention research and is critical in establishing efficacy that can be generalized to a larger population. While this approach also has merit in the context of behavioral interventions, it may unintentionally reduce our ability to detect an effect of the intervention or to differentiate it from an AC (see also²⁶). The efficacy of any legitimate behavioral intervention is predicated upon an individual's engagement with the training and persistence of practice, unlike in a pharmaceutical trial, where one can be fairly confident that every individual is receiving roughly the same dose. Outside the laboratory, individuals choose pathways of change that they are most drawn to, or for which they have some aptitude. Choice is a strong predictor of adherence to and engagement with an intervention^{27,28} and effect sizes are typically higher when an intervention is individually initiated, rather than the consequence of random assignment²⁹. Thus, through random assignment, these studies may unintentionally deflate the effect size of the intervention and the question must be posed: to which

population is it most important to generalize — to a truly random sample or to a population of individuals who would initiate participation in an MBSR intervention? In order to estimate an effect size that would be comparable to those of MBSR interventions offered in the community, it would be necessary to allow participants to choose between MBSR and AC interventions. To determine the difference in effect size for outcomes of MBSR interventions between self-initiated and random assignment, it would be necessary to randomly assign participants to choice or random assignment. While theoretically compelling, the latter suggestion is not feasible under most current funding mechanisms and the former is vulnerable to biases associated with self-selection. This remains an important and challenging issue that has been largely unexamined. At minimum, participant preference should be recorded prior to random assignment and modeled in subsequent analyses, including as a potential confounder, mediator or moderator of important outcome measures.

The question to whom should we aim to generalize can be extended to sample selection as well. Sample selection is a crucial factor in establishing an effect of the intervention. On one hand, restrictions to inclusion are necessary in order to avoid confounds in key outcome measures. In addition, limiting sample heterogeneity with precise inclusion criteria reduces variance in baseline measures. On the other hand, the selected sample should be responsive to the intervention and resemble the population to which it is meant to generalize. As a field, we may put too much weight on the former at the expense of the latter. In the wild, many individuals seek out MBSR in search of relief from suffering in

some form. Unless the focus of the investigation is a specific clinical sample, these forms of suffering often comprise exclusion criteria in randomized controlled designs, and consequently eliminate from examination those who are most likely to initiate training on their own and perhaps those likely to benefit most from the training. For example, in studies that examine the effect of MBSR on neural function, individuals on psychoactive drugs (e.g. antidepressants or medications for ADHD) are typically excluded. Further, strict sample selection can also result in the creation of a “super normal” set of participants that are functioning exceptionally well, that does not resemble the general population, to which most studies aim to generalize. This may create a ceiling effect for common outcome measures. In addition, motivation to engage with the training and persist at practice once the formal training period is over is typically greatest in those in need of the most relief. By excluding these individuals in a highly selected sample, we may be further reducing the likelihood of differentiating MBSR from ACs by specifically including individuals with a lower likelihood of optimal engagement. To address this issue, the research community should consider broader inclusion criteria, perhaps chosen to reflect the characteristics of those self-enrolling in MBSR courses in the community. Additionally, retrospective data should be collected concerning prior and current adversity, in order to determine if there are optimal conditions in which MBSR and other forms of meditation training are most beneficial.

Expectations regarding the trajectory of change are another factor that complicates the discernment of benefits provided by MBSR training. Inherent in

the methods typically used in analyzing data, it is often assumed that outcome measures should improve in a somewhat linear way, where duration of training or time spent practicing is linearly related to improvement in the outcome measures of interest. This assumption is worth revisiting. It is not uncommon, or unexpected even, for participants to encounter significant difficulties over the course of a mindfulness intervention such as MBSR or MBCT. Segal and colleagues note, for example, that as individuals engage with home practice, they may experience violations of their expectations, such that “They hoped that these practices might make them feel better, yet they may feel worse”³⁰. In a qualitative study of diary entries over the length of an MBSR course, Kerr and colleagues³¹ found that distressful experiences specifically connected to mindfulness practice were common in their participants, even though the negative valence of these experiences did not have an impact on positive outcomes, especially the development of meta-awareness. Finally, traditional sources in Buddhist mindfulness traditions acknowledge that distressful experiences may occur for contemplative practitioners, even though these experiences need not pose insurmountable obstacles for advancement in practice^{32,33}. Depending on the outcome measures deployed in an intervention study, distressful experiences may lead to counterintuitive results, and the expectation of linear improvement in a particular measure may thus be unfounded. Unfortunately, trajectories of change can look very different between individuals, which makes predicting outcomes difficult when just a few snapshots in time are assessed. Individuals who are functioning more poorly at baseline, for

instance may show a larger or temporally-shifted response to MBSR. On the other hand, an individual who is functioning relatively well at baseline will take much more training to realize the same amount of improvement. This pattern, where baseline characteristics moderate intervention response, has been shown with both distress tolerance³⁴ and neuroticism³⁵. Similarly, the trajectory of change may be different for different types of outcome measures. Outcomes that assess changes in “view” for example, may change after very brief training durations³⁶, whereas other outcomes, especially those that require neuroplasticity, may take much more training and practice to show meaningful change^{37,38}. For these reasons, longitudinal studies with periodic sampling would be very valuable. In addition, gathering information that can provide context to the trajectory of change in laboratory measures from participants and close others (e.g. partners, family members, meditation teachers) can provide another lens through which to assess progress.

Last, and perhaps most important, is the intention with which participants engage with mindfulness training. One of the starker differences between training in meditation and training contained in an AC is the focus of the benefits that the training will yield. Though frequently viewed by the public as a self-help strategy, the style of mindfulness cultivated in MBSR is intended to enhance one’s connections with others, especially in relation to universally shared goals of alleviating suffering and enhancing life. The notion that mindfulness in MBSR should prompt an other-centered perspective in this way was an explicit element in the development of the program, at least according to its primary architect, Jon

Kabat-Zinn^{39,40}. Indeed, in *Full Catastrophe Living*, a foundational publication for MBSR training, Kabat-Zinn speaks of the sense of disconnectedness and fragmentation that often accompanies suffering, and he encourages practitioners to “achieve liberation from the prison of our own thought habits” and its excessive self-focus “intentionally cultivating compassion for all life” (Kabat-Zinn 2005, p.166). But in either case, MBSR could not be construed as a “self-help” program focused on benefitting only each individual participant. In contrast, in most cases the intention for the benefits of ACs are entirely self-focused, at least in a direct sense.

As within any training program, there will be variability in the degree to which MBSR instruction actually conveys and realizes the paradigmatic processes and goals of the program, including a shift away from a self-centered perspective to one that is more other-centered. When MBSR training does create such a shift, an enhanced sense of social connection and a less rigid sense of self are likely to result through mechanisms that differentiate MBSR from ACs that do not target such a shift. For example, frequency of self-related thought is positively associated with symptoms of anxiety and depression⁴¹ and is reflected in altered activity and functional connectivity in the default mode network^{42,43}. In these two domains, MBSR training, at least in its paradigmatic form, should produce changes that will lead MBSR to be more differentiated from ACs in measures of frequency of self-related thought and DMN activity and connectivity (e.g.^{44,45}, whereas symptoms of anxiety and depression alone may fail to differentiate the interventions. An important caveat here, however, is the degree

to which self-focus may be an especially difficult target of change in cultures such as the United States, where a highly individualistic approach to identity is so dominant⁴⁶. In any given MBSR study, it thus may be especially important to assess treatment fidelity in regard to this aspect of the MBSR protocol.

The application of modern scientific inquiry and tools to better understand the nature and benefits of contemplative practices like meditation is in its infancy. Nonetheless, an enormous amount of data have been generated already and the rate of new publications on this topic is growing nearly exponentially⁴⁷. We are now in a position to take a close and careful look at what has been learned so far and use this opportunity to refine our experimental questions and approaches to address more nuanced hypotheses. The employment of well-matched ACs in the investigation of the benefits of MBSR brings us one step closer to the level of rigor necessary to establish meditation practices as efficacious and efficient ways to promote wellbeing and reduce suffering for years to come. Yet more work is needed to delineate how, when, why, and for whom these practices are preferable to other available options.

Conflicts of Interest

Dr. Richard J. Davidson is the founder, president, and serves on the board of directors for the non-profit organization, Healthy Minds Innovations, Inc. In addition, Dr. Davidson served on the board of directors for the Mind & Life Institute from 1992-2017. Dr. Rosenkranz and Prof. Dunne have no conflicts of interest to disclose.

Acknowledgments

This research was supported by the National Center for Complementary and Integrative Health (NCCIH) P01AT004952 to RJD & MAR, the National Heart, Lung, and Blood Institute (NHLBI) R01HL123284 to William W. Busse, and generous donations from individuals to the Center for Investigating Healthy Minds.

References Cited

1. Khoury, B., Sharma, M., Rush, S. E. & Fournier, C. Mindfulness-based stress reduction for healthy individuals: A meta-analysis. *J. Psychosom. Res.* **78**, 519–28 (2015).
2. Gotink, R. A. *et al.* Standardised mindfulness-based interventions in healthcare: An overview of systematic reviews and meta-analyses of RCTs. *PLoS One* **10**, 1–17 (2015).
3. Fjorback, L. O., Arendt, M., Ornbol, E., Fink, P. & Walach, H. Mindfulness-based stress reduction and mindfulness-based cognitive therapy - a systematic review of randomized controlled trials. *Acta Psychiatr. Scand.* **124**, 102–119 (2011).
4. Lakhan, S. E. & Schofield, K. L. Mindfulness-Based Therapies in the Treatment of Somatization Disorders: A Systematic Review and Meta-Analysis. *PLoS One* **8**, (2013).
5. Katterman, S. N., Kleinman, B. M., Hood, M. M., Nackers, L. M. & Corsica, J. A. Mindfulness meditation as an intervention for binge eating, emotional eating, and weight loss: A systematic review. *Eat. Behav.* **15**, 197–204 (2014).
6. O'Leary, K., O'Neill, S. & Dockray, S. A systematic review of the effects of mindfulness interventions on cortisol. *J. Health Psychol.* 1359105315569095- (2015). doi:10.1177/1359105315569095
7. Goyal, M. *et al.* Meditation programs for psychological stress and well-

- being. *JAMA Intern. Med.* **174**, 357 (2014).
8. Coronado-Montoya, S. *et al.* Reporting of positive results in randomized controlled trials of mindfulness-based mental health interventions. *PLoS One* **11**, 1–18 (2016).
 9. MacCoon, D. G. *et al.* The validation of an active control intervention for Mindfulness Based Stress Reduction (MBSR). *Behav. Res. Ther.* **50**, 3–12 (2012).
 10. Laska, K. M., Gurman, A. S. & Wampold, B. E. Expanding the Lens of Evidence-Based Practice in Psychotherapy : A Common Factors Perspective. *Psychotherapy* **51**, 467–481 (2014).
 11. Leuchter, A. F., Hunter, A. M., Tartter, M. & Cook, I. A. Role of pill-taking , expectation and therapeutic alliance in the placebo response in clinical trials for major depression. *Br. J. Psychiatry* **205**, 443–449 (2014).
 12. Nyklíček, I. & Kuypers, K. F. Effects of mindfulness-based stress reduction intervention on psychological well-being and quality of life: Is increased mindfulness indeed the mechanism? *Ann. Behav. Med.* **35**, 331–340 (2008).
 13. Hoffman, C. J. *et al.* Effectiveness of mindfulness-based stress reduction in mood, breast- and endocrine-related quality of life, and well-being in stage 0 to III breast cancer: A randomized, controlled trial. *J. Clin. Oncol.* **30**, 1335–1342 (2012).
 14. Tozzi, L. *et al.* Longitudinal functional connectivity changes correlate with mood improvement after regular exercise in a dose-dependent fashion.

- Eur. J. Neurosci.* **43**, 1089–1096 (2016).
15. Black, D. S., O'Reilly, G. a., Olmstead, R., Breen, E. C. & Irwin, M. R. Mindfulness Meditation and Improvement in Sleep Quality and Daytime Impairment Among Older Adults With Sleep Disturbances. *JAMA Intern. Med.* **9239**, 494–501 (2015).
16. Hartescu, I., Morgan, K. & Stevenson, C. D. Increased physical activity improves sleep and mood outcomes in inactive people with insomnia: A randomized controlled trial. *J. Sleep Res.* **24**, 526–534 (2015).
17. Gallegos, A. M., Moynihan, J. & Pigeon, W. R. A Secondary Analysis of Sleep Quality Changes in Older Adults From a Randomized Trial of an MBSR Program. *J. Appl. Gerontol.* (2016).
doi:10.1177/0733464816663553
18. Lutz, A., Jha, A., Dunne, J. D. & Saron, C. D. Investigating the Phenomenological Matrix of Mindfulness-related Practices from a Neurocognitive Perspective. *Am. Psychol.* **70**, 632–658 (2015).
19. Santos, R. V. T., Tufik, S. & De Mello, M. T. Exercise, sleep and cytokines: Is there a relation? *Sleep Med. Rev.* **11**, 231–239 (2007).
20. Youngstedt, S. D. Effects of exercise on sleep. *Clin. Sports Med.* **24**, 355–365 (2005).
21. MacCoon, D. G., MacLean, K. A., Davidson, R. J., Saron, C. D. & Lutz, A. No sustained attention differences in a longitudinal randomized trial comparing mindfulness based stress reduction versus active control. *PLoS One* **9**, (2014).

22. Toubekis, A. G., Tsami, A. P., Smilios, I. G., Douda, H. T. & Tokmakidis, S. P. Training-induced changes on blood lactate profile and critical velocity in young swimmers. *J. Strength Cond. Res.* **25**, 1563–70 (2011).
23. Messonnier, L. A. *et al.* Lactate kinetics at the lactate threshold in trained and untrained men. *J. Appl. Physiol.* **114**, 1593–1602 (2013).
24. Lutz, a, Slagter, H. a, Dunne, J. D. & Davidson, R. J. Attention regulation and monitoring in meditation. *Trends Cogn. Sci.* **12**, 163–169 (2008).
25. Levinson, D. B., Stoll, E. L., Kindy, S. D., Merry, H. L. & Davidson, R. J. A mind you can count on: Validating breath counting as a behavioral measure of mindfulness. *Front. Psychol.* **5**, 1–10 (2014).
26. Davidson, R. J. & Kaszniak, A. W. Conceptual and Methodological Issues in Research on Age and Aging. *Am. Psychol.* **70**, 581–592 (2015).
27. Rennie, T. W., Bothamley, G. H., Engova, D. & Bates, I. P. Patient choice promotes adherence in preventive treatment for latent tuberculosis. *Eur. Respir. J.* **30**, 728–735 (2007).
28. Lindhjem, O., Bennett, C. B., Trentacosta, C. J. & McLear, C. Client preferences affect treatment satisfaction, completion, and clinical outcome: a meta-analysis. *Clin. Psychol. Rev.* **34**, 506–17 (2014).
29. Brown, J. G., Joyce, K. E., Stacey, D. & Thomson, R. G. Patients or volunteers? The impact of motivation for trial participation on the efficacy of patient decision AIDS: a secondary analysis of a Cochrane systematic review. *Med. Decis. Making* **35**, 419–435 (2015).
30. Segal, Z. V, Williams, J. M. G. & Teasdale, J. D. *Mindfulness-based*

- cognitive therapy for depression: A new approach to preventing relapse.* (Guilford Press, 2013).
31. Kerr, C. E., Josyula, K. & Littenberg, R. Developing an observing attitude: an analysis of meditation diaries in an MBSR clinical trial. *Clin. Psychol. Psychother.* **18**, 80–93 (2011).
 32. *Mahamudra, the ocean of true meaning the profound instructions on coexistent unity, the essence of the ocean of true meaning, and light radiating activity.* (Verl.-Haus Monsenstein und Vannerdat, 2009).
 33. Lindahl, J. R., Fisher, N. E., Cooper, D. J., Rosen, R. K. & Britton, W. B. The varieties of contemplative experience: A mixed-methods study of meditation-related challenges in Western Buddhists. *PLoS One* **12**, e0176239 (2017).
 34. Gawrysiak, M. J. et al. Dimensions of distress tolerance and the moderating effects on mindfulness-based stress reduction. *Anxiety, Stress Coping* **29**, 552–560 (2016).
 35. Nyklíček, I. & Irrmischer, M. For Whom Does Mindfulness-Based Stress Reduction Work? Moderating Effects of Personality. *Mindfulness (N. Y.)* **8**, 1106–1116 (2017).
 36. Baquedano, C. et al. Compared to self-immersion, mindful attention reduces salivation and automatic food bias. *Sci. Rep.* **7**, 1–11 (2017).
 37. Hasenkamp, W. & Barsalou, L. W. Effects of Meditation Experience on Functional Connectivity of Distributed Brain Networks. *Front. Hum. Neurosci.* **6**, 1–14 (2012).

38. Allen, M. *et al.* Cognitive-Affective Neural Plasticity following Active-Controlled Mindfulness Intervention. *J. Neurosci.* **32**, 15601–15610 (2012).
39. Kabat-Zinn, J. Some reflections on the origins of MBSR, skillful means, and the trouble with maps. *Contemp. Buddhism* **12**, 281–306 (2011).
40. Kabat-Zinn, J. Too Early to Tell: The Potential Impact and Challenges-Ethical and Otherwise-Inherent in the Mainstreaming of Dharma in an Increasingly Dystopian World. *Mindfulness (N. Y.)*. **8**, 1125–1135 (2017).
41. Hoffmann, F., Banzhaf, C., Kanske, P., Bermpohl, F. & Singer, T. Where the depressed mind wanders: Self-generated thought patterns as assessed through experience sampling as a state marker of depression. *J. Affect. Disord.* **198**, 127–134 (2016).
42. Nejad, A. B., Fossati, P. & Lemogne, C. Self-Referential Processing, Rumination, and Cortical Midline Structures in Major Depression. *Front. Hum. Neurosci.* **7**, 666 (2013).
43. Kaiser, R. H. *et al.* Dynamic Resting-State Functional Connectivity in Major Depression. *Neuropsychopharmacology* **41**, 1–9 (2015).
44. Creswell, J. D. *et al.* Alterations in resting state functional connectivity link mindfulness meditation with reduced interleukin-6: a randomized controlled trial. *Biol. Psychiatry* 1–9 (2016). doi:10.1016/j.biopsych.2016.01.008
45. Garrison, K. A., Zeffiro, T. A., Scheinost, D., Constable, T. & Brewer, J. A. Meditation leads to reduced default mode network activity beyond an active task. *Cogn. Affect. Behav. Neurosci.* **15**, 712–720 (2015).
46. Triandis, H. C., Bontempo, R., Villareal, M. J., Asai, M. & Lucca, N.

- Individualism and collectivism: Cross-cultural perspectives on self-ingroup relationships. *J. Pers. Soc. Psychol.* **54**, 323–338 (1988).
47. Black, D. S. Mindfulness-based interventions: an antidote to suffering in the context of substance use, misuse, and addiction. *Subst. Use Misuse* **49**, 487–91 (2014).